

Influenza (Flu) Information for Parents

For parents

Should my child get a flu vaccination this year?

Yes. Everyone older than 6 months of age should get flu vaccine this year.

While everyone should be vaccinated, it is especially important for children at risk for serious complications of the flu. These include:

- Children age 6 months to 5 years
- Those age 5-18 years with:
 - Heart disease
 - Pulmonary disease including asthma
 - Metabolic disease (diabetes)
 - Immune deficiency
 - Blood disorders
 - Long-term aspirin therapy
- Teens who are pregnant during the influenza season

Getting a flu shot (or nasal spray) means your child won't catch the flu, so he or she won't pass it to people who risk getting very sick – like babies, elderly people, and people who have chronic diseases.

Should my child get one or two doses of flu vaccine?

That will depend on your child's age and whether they got flu vaccine this past year. If your child is 6 months through 8 years old, he or she may need two doses, at least four weeks apart. Ask your doctor or clinic.

Are we still in an H1N1 flu pandemic?

No, but we expect the H1N1 flu strain to continue to circulate and cause disease this season. That's why the H1N1 strain of the virus is included in this year's seasonal flu vaccine.

I heard that this year's flu vaccine has H1N1 in it. Is it safe?

Yes. This year's flu vaccine is made in the same way as past flu vaccines. Flu vaccines have an excellent safety record – with an average of 100 million doses used in the U.S. each year.



Immunization Program
 P.O. Box 64975
 St. Paul, MN 55164-0975
 651-201-5503, 1-800-657-3970
www.mdhflu.com

Are children at high risk for complications of influenza?

Most healthy children can weather the flu, even with a high fever. But that's not always true for children younger than 5 years old and those with chronic conditions like asthma and diabetes.

Serious complications of flu occur most often in young infants, people with chronic health conditions, and the elderly.

Sometimes healthy people – including children – can have these serious complications or die from the flu. That's why we need to take flu very seriously and get vaccinated each year.

Everyone older than 6 months of age should get flu vaccine this year.

What is influenza (flu)?

Influenza is a contagious respiratory disease that can be prevented by immunization. It is not the same as the "stomach flu." Flu is caused by a virus that attacks the nose, throat, and lungs. It can cause mild to severe illness and at times can lead to death.

What are the symptoms of influenza?

Influenza symptoms come on quickly in the form of fever, dry cough, sore throat, headache, extreme tiredness, stuffed-up nose, and body aches. These symptoms can be severe and put you in bed for several days.

How is flu different from a cold?

A cold generally stays up in the head while the flu brings body aches, fever, and extreme fatigue. A child with a cold will usually keep up with normal activities. A child with the flu will often feel too sick to play.

What about the nasal vaccine, FluMist?

Healthy individuals 2 through 49 years of age, including those who have contact with or are caregivers of infants under 6 months of age, can receive FluMist. Check with your doctor or clinic.

Influenza (Flu) Information for Parents – page 2

What if my child already had H1N1 flu disease last season?

Even if your child had H1N1 flu, other strains of flu virus that are included in the vaccine will be circulating this season, and it is important to be protected from those as well.

My child just got flu vaccine last spring. Is it too soon to get another flu shot now?

No, it's not too soon. Other strains of flu virus that are included in this season's vaccine will be circulating, and it is important to be protected from those as well.

What if you think you or your child has the flu?

- Stay home if you are ill and keep your child home from school or daycare if they are ill.
- Rest and drink lots of fluids.
- Antibiotics will not help a person recover from the flu, because flu is caused by a virus, not by bacteria.
- Children often need help keeping their fever under control. Follow your child's doctor's instructions.
- Take your child to the doctor or the emergency room if he or she:
 - Breathes rapidly or with difficulty
 - Has bluish skin color
 - Does not drink enough and becomes dehydrated
 - Does not wake up or interact with others
 - Is so irritable that he or she doesn't want to be held, or
 - Gets better only to become sick again, with fever and a more severe cough
- If you are concerned that something does not seem right with your child, call your doctor or clinic.

What can you do to protect yourself and others?

- Get vaccinated.
- Avoid being exposed to others who are sick with a flu-like illness.
- Cover your nose and mouth with a tissue when you cough or sneeze, or cough or sneeze into your sleeve.

- Clean your hands often – with soap and water or a hand sanitizer.
- Don't expose infants unnecessarily to large crowds when influenza is in your community. Avoid close contact between the baby and family members who may be sick.
- Do not share drinking cups and straws.
- Frequently clean commonly touched surfaces (door knobs, refrigerator handles, phones, water faucets).
- Do not smoke around children.

What about antiviral medicines?

Antiviral medicines can offer some protection, but they are only recommended for use in certain groups of people. If you have questions about antivirals, talk to your doctor.

**Find a flu shot at
www.mdhflu.com**